arXiv:gr-qc/0606004v2 20 Jun 2006

Selfgravitation in a general-relativistic accretion of steady fluids

Bogusz Kinasiewicz,[1] Patryk Mach,[1] and Edward Malec[1]

[1]*M. Smoluchowski Institute of Physics, Jagiellonian University, Reymonta 4, 30-059 Kraków, Poland*

The selfgravity of an infalling gas can alter significantly the accretion of gases. In the case of spherically symmetric steady flows of polytropic perfect fluids the mass accretion rate achieves maximal value when the mass of the fluid is 1/3 of the total mass. There are two weakly accreting regimes, one over-abundant and the other poor in fluid content. The analysis within the newtonian gravity suggests that selfgravitating fluids can be unstable, in contrast to the accretion of test fluids.

INTRODUCTION.

The analysis of the spherical steady accretion of gas onto a gravitational center starts with the Newtonian accretion discussed by Bondi [1]. Michel [2], Shapiro and Teukolsky [3], and others [4] investigated the accretion of test fluids in the Schwarzschild space-time. Recent analysis showing the importance of backreaction is that of Karkowski et al. [5]. In this lecture we follow the general-relativistic description of selfgravitating fluids that has been formulated in [6] and present main results of [5].

One of the reasons for inspecting steady flows in the full generality, when infalling gas modifies the space-time geometry, is the wish to see the effects of backreaction in a simple but nontrivial accretion model. We show later, following [5], that the mass accretion rate $\dot{m}$ behaves like $(1-x)^2\dot{m}_B$, where $x = m_f/m$ is the ratio of the mass of the fluid to the total mass and $\dot{m}_B$ is the mass accretion rate for the accretion of test fluids. Since $\dot{m}_B$ is proportional to the fluid abundance, hence to x, one sees that there exists a maximum of $\dot{m}$ when $x = 1/3$. The mass accretion rate $\dot{m}$ sets an upper limit for the luminosity of the system. This simple analysis of accretion suggests the existence of two weakly luminous regimes: one rich in fluid with $m_f \approx m$ and the other with a small amount of fluid, $m_f/m \ll 1$. It is clear from this reasoning that it is reasonable to take into account selfgravitation of accreting gases whenever their mass is comparable to the central mass. On the other hand, the stability properties of accreting fluids can depend on backreaction, however small. The stability of steady flows in a fixed relativistic background as well as in the newtonian context has been shown by Moncrief [7]. We show later that the steady newtonian acretion of selfgravitating gases does not exclude instability.

Trying to put matter in the astrophysical context, notice that one can expect the mass of accreting fluids to be comparable to the central mass only in the case of heavy and luminous compact objects; compact galactic centers probably constitute fair examples. This suggests that the accretion physics of such objects should include the selfgravity of accreting matter. Let us mention here that there exists a fundamental physical aspect that the astrophysics of accreting matter might be able to resolve. Namely, the standard point of view is that compact objects of a mass exceeding several solar masses constitute black holes. That interpretation has been challenged by Mazur and Mottola [8] who developed the notion of gravatars — stars almost as compact as black holes but dispossessed of event horizons. Gravastars are supposed to be built from matter violating energy conditions [9] and in this way they would avoid the standard compactness and mass limitations, like the Buchdahl theorem [10] or Chandrasekhar type bounds on mass. There is an interesting motivation coming from the quantum-gravity type arguments, that inspired the concept of gravastars, but we leave it aside. It is disputable whether observations can distinguish compact bodies endowed with a hard surface from black holes within the known nowadays accretion models. For a discussion of that issue see [11] and [12]. We believe that the resolution of that problem requires the inclusion of general-relativistic aspects such as backreaction. In this talk we shall report the recent progress in this direction.

MAIN EQUATIONS.

We will consider a spherically symmetric compact ball of a fluid falling onto a non-rotating black hole. The equations are taken from [6]. We will use comoving coordinates

$$ds^2 = -N^2 dt^2 + \alpha dr^2 + R^2\left(d\theta^2 + \sin^2\theta d\phi^2\right), \qquad (1)$$

where the lapse N, α and the areal radius R are functions of a coordinate radius r and an asymptotic time variable t. The nonzero components of the extrinsic curvature K_{ij} of the t = const slices read $K^r_r = \frac{1}{2N\alpha}\partial_t\alpha$, $K^\phi_\phi = K^\theta_\theta = \frac{\partial_t R}{NR} = \frac{1}{2}(\mathrm{tr}K - K^r_r)$. Here $\mathrm{tr}K = \frac{1}{N}\partial_t \ln\left(\sqrt{\alpha}R^2\right)$. The mean curvature of two-spheres of constant radius r, embedded in a Cauchy hypersurface is $k = \frac{2\partial_r R}{R\sqrt{\alpha}}$. The energy-momentum tensor of the perfect fluid reads $T^{\mu\nu} = (p+\varrho)u^\mu u^\nu + pg^{\mu\nu}$ (where we assume the barotropic equation of state), $u_\mu u^\mu = -1$ with u^μ denoting the four-velocity of the fluid, p is the pressure and ϱ the energy density in the comoving frame.

The areal velocity U of a comoving particle labelled by coordinates (r,t) is given by $U(r,t) = \frac{\partial_t R}{N} = \frac{R}{2}(\mathrm{tr}K - K^r_r)$. From the Einstein constraint equations one has

$$kR = 2\sqrt{1 - \frac{2m(R)}{R} + U^2}. \qquad (2)$$

Here and below we use $\partial_r = \sqrt{\alpha}(kR/2)\partial_R$ in order to eliminate the differentiation with respect to the comoving radius r. The quasilocal mass $m(R)$ is defined by $\partial_R m(R) =$

$4\pi R^2\varrho$. The mass accretion rate is $\dot{m} = (\partial_t - (\partial_t R)\partial_R)m(R) = -4\pi N R^2 U(\varrho + p)$ or (equivalently, but perhaps more familiarly) $\dot{m} = -4\pi R^2 nU$, where n is the baryonic density. We assume a steady collapse of the fluid, which means that all its characteristics are constant at a fixed areal radius R: $\partial_t X|_{R=\mathrm{const}} = (\partial_t - (\partial_t R)\partial_R)X = 0$, where $X = \varrho, U, a^2$. $a = \sqrt{\partial_\varrho p}$ is the speed of sound. Strictly saying, a stationary accretion must lead to to the increase of the central mass and of some geometric quantities. This in turn means that the notion of the "steady accretion" is approximate — it demands that the mass accretion rate is small and the time scale is short, so that the quasilocal mass $m(R)$ does not change significantly. Thence also geometric quantities, like the mean curvature k or the area of the black hole, remain practically constant.

The Einstein evolution equation $\partial_t U = k^2R^2\partial_R N/4 - m(R)N/R^2 - 4\pi NRp$ and the energy conservation equation $\partial_t\varrho = -N\mathrm{tr}K(\varrho + p)$ constitute (assuming steady flow) ordinary differential equations with respect to R.

It is easy to show that now the mass accretion rate is constant, $\partial_R\dot{m} = 0$ [6].

By a black hole is understood a region within an apparent horizon to the future, i.e., a region enclosed by an outermost sphere S_A on which the optical scalar $\theta_+ \equiv \frac{pR}{2} + U$ vanishes [14]. (The other condition, that $\theta_- \equiv \frac{pR}{2} - U > 0$ for all spheres outside S_A, is satisfied trivially for a steady accretion.) That means that on S_A the ratio $2m_{BH}/R_{AH}$ becomes 1, where R_{AH} is the areal radius of the apparent horizon and $m_{BH} \equiv m(R_{AH})$ is the mass of the black hole. We assume the polytropic equation of state $p = K\varrho^\Gamma$, with Γ being a constant, $(1 < \Gamma \leq 5/3)$.

Furthermore, we assume that the radius R_∞ of the ball of fluid and other boundary data are such $|U_\infty| \ll \frac{m(R_\infty)}{R_\infty} \ll a_\infty$. These boundary conditions allow one to join smoothly the internal solution (the steady fluid) with the external Schwarzschild geometry. The momentum conservation equation $\nabla_\mu T^\mu_r = 0$ — in comoving coordinates

$$N\partial_R p + (p + \varrho)\,\partial_R N = 0 \tag{3}$$

— can be integrated, yielding (with $N(R_\infty) = 1$)

$$a^2 = -\Gamma + \frac{\Gamma + a^2_\infty}{N^\kappa}. \tag{4}$$

Here $\kappa = \frac{\Gamma-1}{\Gamma}$. Equation (4) is the general-relativistic version of the Bernoulli equation. Equation (3) implies also that the lapse expresses as follows $N = \frac{Cn}{p+\varrho}$, with the baryonic number density $n = C\exp\int_{\varrho_\infty}^{\varrho} ds\frac{1}{s+p(s)}$. The baryonic current is conserved, $\nabla_\mu\left(nu^\mu\right) = 0$.

The sonic point is as a location where the length of the spatial velocity vector $|\vec{U}| = |U|/(kR/2)$ equals a. Therefore $|U| = \frac{1}{2}kRa$. In the Newtonian limit this coincides with the standard requirement $|U| = a$. In the following we will denote by the asterisk all values referring to the sonic points, e.g. a_*, U_*, etc. The four characteristics, a_*, U_*, m_*/R_* and c_* are related [6], $a^2_*\left(1 - \frac{3m_*}{2R_*} + c_*\right) = U^2_* = \frac{m_*}{2R_*} + c_*$, where $c_* = 2\pi R^2_* p_*$. The infall velocity U reads

$$U = U_*\frac{R^2_*}{R^2}\left(\frac{1 + \frac{\Gamma}{a^2}}{1 + \frac{\Gamma}{a^2_*}}\right)^{1/(\Gamma-1)}. \tag{5}$$

Here U_* is the negative square root. From the relation between the pressure and the energy density, one obtains, using equation (4)

$$\varrho = \varrho_\infty\left(a/a_\infty\right)^{2/(\Gamma-1)} = \varrho_\infty\left(-\frac{\Gamma}{a^2_\infty} + \frac{\frac{\Gamma}{a^2_\infty} + 1}{N^\kappa}\right)^{\frac{1}{\Gamma-1}}, \tag{6}$$

where the constant ϱ_∞ is equal to the mass density of a collapsing fluid at the boundary R_∞. The steady fluid is described by equations (4 – 7). They constitute an integro-algebraic system of equations, with a bifurcation point at the sonic point, where two branches (identified as an accretion or a wind) do cross; that is a well known feature of that problem, present also in models with a test fluid [1] – [4], [6]. That requires some caution in doing numerics and a careful selection of the solution branch. Notice that these equations are expressed exclusively in terms of quantities that are steady in the sense of the former definition.

CALCULATION OF THE MASS ACCRETION RATE.

In numerical calculations it will be convenient to represent the mass $m(R)$ in the form

$$m(R) = m - 4\pi\int_R^{R_\infty} drr^2\varrho. \tag{7}$$

Notice that $m(\infty) = m_\infty \approx m$. For a steady flow one obtains [6]

$$\begin{aligned} N &= \frac{kR}{k(R_\infty)R_\infty}\beta(R), \\ \beta(R) &= \exp\left(-16\pi\int_R^{R_\infty}(p + \varrho)\,\frac{ds}{k^2 s}\right). \end{aligned} \tag{8}$$

The full description of the steady accretion is given by equations (2), (7), (4), (5) and (8), which are equivalent to the former set of equations.

In the forthcoming part of the talk we shall describe main points of the derivation of the accretion mass rate, following the presentation of Karkowski et al. [5].

In the first step, one shows in [5], that outside the sonic horizon solutions of the equations with backreaction can be estimated by relevant solutions describing the stationary accretion of test fluids. We supplement the foregoing list of conditions by another, that the mass m_f of the fluid outside the sonic horizon is of the order of the mass m_* within the sonic horizon; That allows one to show that the function β appearing in equation (8) is essentially 1 and $c_* = 2\pi R^2_* p_* \ll a^2_*$ can be ignored. As a consequence one infers that in the case of stationary accretion of selfgravitating fluids some characteristics of the sonic point — a_*^2, U_*^2 and m_*/R_* — are the same as for the test fluid accretion, provided that asymptotic data ρ_∞ and a_∞^2 are identical [5]. In particular, in both cases characteristics of the sonic point do not depend on the mass — either the total mass m or the mass m_* of the accretor.

In the second step one shows that it is possible to estimate the energy density ϱ of a system with backreaction using information about solutions in the test fluid approximation. It is done in [5]. We shall describe in a more detail how one obtains the desired form of estimates onto the energy density of test fluids. One starts with the estimate on the inverse of the lapse function $N(\delta) = \sqrt{1-\delta}$, where $\delta \equiv \frac{2m}{R} - U^2$ is less than one (and in fact strictly smaller than 3/4). It is easy to show, applying the method of differential inequalities, that $N^{-1}(\delta) \leq 1 + \delta/\left(2N^3(\delta)\right)$. Since $\delta \leq 2m/R$, one has $N^{-1}(\delta) \leq 1 + m/\left(RN^3(\delta)\right)$. Notice now that the lapse is an increasing function and $N(R) \geq N_* \equiv N(R_*)$ for $R \geq R_*$. Therefore one arrives at the inequality

$$N^{-1} \leq 1 + \frac{1}{N_*^3} \times \frac{m}{R}. \tag{9}$$

Having that, one can derive a bound onto ϱ. One obtains, applying the inequality (9),

$$\varrho(R) \leq \varrho_\infty \left(1 + (\Gamma - 1)\frac{1 + \frac{\Gamma}{a_\infty^2}}{\Gamma N_*^3} \times \frac{m}{R}\right)^{1/(\Gamma-1)}. \tag{10}$$

The lapse at the sonic point $N_* = 1/\sqrt{1+3a_*^2}$ and $m/R_* = a_*^2 N_*^2$; thus a constant $\alpha \equiv \frac{1+\frac{\Gamma}{a_\infty^2}}{\Gamma N_*^3}\frac{m}{R_*}$ can be written as

$$\alpha = \frac{1}{\Gamma}\sqrt{1+3a_*^2}\left(a_*^2 + \frac{\Gamma a_*^2}{a_\infty^2}\right). \tag{11}$$

One can prove following bounds (see equation (5.2) in [6])

$$\frac{a_*^2}{a_\infty^2} \leq \frac{2}{5 - 3\Gamma + \frac{3a_*^2(\Gamma-1)^2(9\Gamma-7)}{4\Gamma\left(1+3a_*^2\right)}}. \tag{12}$$

This implies a series of estimates: i) for $1 < \Gamma \leq 4/3$ one has $\frac{a_*^2}{a_\infty^2} < 2$; ii) in the range $4/3 < \Gamma \leq 3/2$ holds $\frac{a_*^2}{a_\infty^2} < 4$; iii) when $\Gamma > 3/2$ and $a_*^2 < 0.2$ one obtains $\frac{a_*^2}{a_\infty^2} < 2.3/a_*^2$ and $a_*^2 \leq 1.6a_\infty$, iv) while for $\Gamma > 3/2$ and $a_*^2 \geq 0.2$ it follows $\frac{a_*^2}{a_\infty^2} < 4$. Now, if we demand that the sonic point is outside of an apparent horizon then $m/(2R_*) < 1/4$ and that implies $a_*^2 \leq 1$. Using this fact, one can derive bounds onto the constant α. Namely, in each of the above cases we have i) $1 < \Gamma \leq 4/3 \to \alpha < 6$; ii) $4/3 < \Gamma \leq 3/2 \to \alpha < 9.5$; iii) $\Gamma > 3/2$ and $a_*^2 < 0.2$: $\alpha < 1 + 2/a_\infty$ iv) $\Gamma > 3/2$ and $a_*^2 \geq 0.2$: $\alpha < 8.3$.

With these estimates one obtains simple bounds onto ρ. We shall consider in more detail two of them corresponding to the cases i) and iii). For $\Gamma \leq 4/3$ equation (10) can be written, using the inequality $(1 + A/x)^x \leq e^A$ and $\alpha \leq 6$, as

$$\varrho(R) \leq \varrho_\infty \exp\left(\frac{6R_*}{R}\right). \tag{13}$$

In the case iii) one notices that $(1+A)^{\frac{1}{\Gamma-1}} \leq 1 + 2A + A^2$. That leads to the estimate

$$\frac{\varrho}{\varrho_\infty} \leq 1 + 2(\Gamma-1)\alpha\frac{R_*}{R} + (\Gamma-1)^2\alpha^2\frac{R_*^2}{R^2}. \tag{14}$$

one can check the selfconsistency of the calculation — for instance that the function $\beta(R) \approx 1$ for $R \geq R_*$. Indeed, the exponent in the exponential funtion $\beta(R_*)$ is of the order of $M_f \equiv 4\pi\int_{R_*}^{R_\infty} drr\rho$. This latter expressions reads in the case i)

$$\begin{aligned} M_f = {} & 2\pi\rho_\infty R_\infty^2 \Bigg[e^{6R_*/R_\infty} - e^6\frac{R_*^2}{R_\infty^2} + \\ & \frac{6R_*}{R_\infty}e^{6R_*/R_\infty} - 6e^6\frac{R_*^3}{R_\infty^3} + \\ & \left(\frac{36R_*^2}{R_\infty^2}e^{\frac{6R_*}{R_\infty}} - 36e^6\frac{R_*^4}{R_\infty^4}\right)\int_1^{R_\infty/(6R_*)} dx\frac{e^{1/x}}{x}\Bigg]. \end{aligned} \tag{15}$$

Since $R_\infty \gg R_*$ one concludes that $M_f \approx 2\pi\rho_\infty R_\infty^2$ which in turn is clearly much smaller than m/R_*. Thus the exponent of $\beta(R)$ is negligibly small for $R \geq R_*$ and $\beta(R) \approx 1$ as desired. A similar reasoning in the case iii) leads to $M_f \approx 2\pi\rho_\infty R_\infty^2\left(1 + O\left(\alpha R_*/R_\infty\right)\right)$. Now $\alpha R_*/R_\infty \leq R_*/R_\infty + 2R_*/\left(R_\infty a_\infty\right)$. It is easy to see, taking into account the boundary condition $a_\infty^2 \gg m/R_\infty$ and $a_* \approx a_\infty$, that the second term in M_f is much smaller than 1 and therefore the expression $O\left(\alpha R_*/R_\infty\right)$ is negligible small. Thus again $M_f \approx m_f/R_\infty \ll m/R_\infty \ll 1$, where m_f is the mass of fluid outside the sonic horizon; therefore one again arrives at $\beta(R_*) \approx 1$.

In the third step one shows — using the same reasoning as in the preceding paragraghs — that the mass $m_f = \gamma\varrho_\infty$, where γ is a constant. This is because the mass density changes moderately outside the sonic point, as shown by estimates (13-14). The calculation is lengthy but simple and we omit it here. Taking into account $m = m_* + m_f$, one infers $m_*/m = 1 - \varrho_\infty(\gamma/m)$; the ratio $x \equiv m_*/m$ linearly decreases with the growth of ϱ_∞.

The rate of the mass accretion $\dot{m}$ can be expressed as below (see equation (6.1) in [6])

$$\dot{m} = \pi m_*^2\varrho_\infty\frac{R_*^2}{m_*^2}\left(\frac{a_*^2}{a_\infty^2}\right)^{\frac{(5-3\Gamma)}{2(\Gamma-1)}}\left(1 + \frac{a_*^2}{\Gamma}\right)\frac{1+3a_*^2}{a_\infty^3}. \tag{16}$$

The whole dependence on ϱ_∞ is contained in the factor $m_*^2\varrho_\infty = m^3x(1-x)^2/\gamma$. Thus one can write

$$\dot{m} = m^3x\frac{(1-x)^2}{\gamma}\pi\frac{R_*^2}{m_*^2}\left(\frac{a_*^2}{a_\infty^2}\right)^{\frac{(5-3\Gamma)}{2(\Gamma-1)}}\left(1 + \frac{a_*^2}{\Gamma}\right)\frac{1+3a_*^2}{a_\infty^3}. \tag{17}$$

This expression clearly demonstrates that the mass accretion rate achieves a maximum at $m_* = 2m/3$ and tends to zero when i) $m_* \to m$ (when the density ϱ tends to zero) and ii) $m_*/m \to 0$.

Thus, one concludes that **Amongst steadily accreting systems of the same R_∞, a_∞ and m those will be most efficient for which $m_* = 2m/3$.** The factor 2/3 is universal — independent of the parameters R_∞, Γ and a_∞.

Since the mass of the fluid m_f is close to $m - m_*$ and $m = m_f + m_{BH}$, the above means that the maximum of the mass accretion takes place when the mass of the fluid is a half of the mass of the black hole.

Finally let us mention that in the case of polytropes with $= Kn^\Gamma$ the Bernoulli equation reads $N\left(\Gamma - 1 - a_\infty^2\right) = \Gamma - 1 - a^2$. The analytic arguments of the type presented above can be applied and conclusions are the same. Similar results should be true for massive fluids in the newtonian limit provided that the polytropic index Γ is clearly smaller than $5/3$.

NUMERICAL RESULTS.

In this Section we present a small sample of the numerical investigation, different from that shown in [5]. The main lesson that one can infer from these is that boundary conditions displayed in Sec. 2 can be significantly weakened. It is clear from considerations around (12) that the point where one can expect difficulties is when the adiabatic index $\Gamma = 5/3$, since in this case the relevant bounds are relatively poor. But the numerical data, that are displayed in the forthcoming Table and figures, demonstrate suprisingly good agreement with the statements of the preceding section.

TABLE I: Characteristics of the sonic point a_*^2, $|U_*|$, R_* for the $\Gamma \approx 5/3$ polytrope and $a_\infty^2 = 0.1$. The last column shows the areal radius of the apparent horizon. The mass accretion rate (second column) achieves maximum at $m_{BH} \approx 0.663$.

m_f/m	$\dot{m}$	R_*	U_*	a_*^2	R_{AH}
4.18×10^{-32}	-2.75×10^{-48}	2.76	0.424584	0.397394	1.99
4.18×10^{-22}	-2.75×10^{-38}	2.76	0.424584	0.397394	1.99
4.18×10^{-12}	-2.75×10^{-28}	2.76	0.424584	0.397394	1.99
0.038	-2.53×10^{-18}	2.64	0.424584	0.397395	1.91
0.075	-4.62×10^{-18}	2.53	0.424621	0.397703	1.82
0.12	-6.31×10^{-18}	2.41	0.424621	0.397702	1.74
0.16	-7.63×10^{-18}	2.30	0.424575	0.397359	1.66
0.2	-8.60×10^{-18}	2.18	0.424621	0.397702	1.57
0.25	-9.26×10^{-18}	2.06	0.424621	0.397703	1.49
0.29	-9.63×10^{-18}	1.95	0.424621	0.397719	1.40
0.33	-9.73×10^{-18}	1.83	0.424619	0.397784	1.32
0.38	-9.62×10^{-18}	1.72	0.424585	0.397395	1.24
0.41	-9.30×10^{-18}	1.60	0.424611	0.397586	1.16
0.83	-1.45×10^{-18}	0.44	0.424621	0.397707	0.32
0.88	-8.37×10^{-19}	0.33	0.424621	0.397707	0.24
0.94	-2.05×10^{-19}	0.16	0.424621	0.397704	0.11

It appears that the characteristics a_*^2, U_*^2, m_*/R_* of the sonic point *practically* do not depend, for a given Γ and a_∞^2, on the energy density ϱ_∞ (exemplary results are shown in Table 1), suggesting a precision of 10^{-5}. The more reliable measure of the numerical error is the difference between the mass $m = 1$ (assumed in the equations) and the mass found numerically. Our results (see Table) suggest that numerical error is smaller than 0.5 %. The quantity c_* is negligible in comparison to other sonic point parameters. The mass m_* behaves like $m_* = 1 - \gamma\varrho_\infty$ (Fig. 1). And finally, the extremum of $\dot{m}$ is achieved when $m_* \approx 2/3$ (Fig. 2). Notably, numerical results extend the regime in which the universality is observed into fluids characterized by the speed of sound a_∞ exceeding 1. This is remarkable, since that implies $R_* < R_{AH}$ and the theoretical analysis presented above would not be valid.



FIG. 1: The dependence of $\frac{m_f}{m} \approx 1 - m_*$ on the asymptotic mass density.



FIG. 2: The dependence of $\dot{m}$ on the ratio $(m - m_*)/m$.

Formula (16) implies that the mass accretion rate in the case with backreaction (when $m_* < m$) can be only smaller from the mass accretion rate without backreaction (when $m_* = m$), with the same asymptotic parameters ρ_∞, a_∞, m. Numerical results agree with the analytic conclusion that the proportionality coefficient γ depends essentially only on the size of support.

STABILITY OF SELFGRAVITATING FLUIDS IN LAGRANGIAN COORDINATES.

The effect of selfgravitation of the steadily accreting gases on the stability has not been studied, up to our knowledge, even in the Newtonian case. In this section we shall fill this gap by analyzing nonrelativistic accretion. The relativistic accretion will be done elsewhere. The forthcoming analysis holds true for polytropes and, more generally, for barotropes. The newtonian equations for selfgravitating fluids in terms of the Eulerian time t and the areal radius R read

$$\begin{aligned} \partial_t U + U\partial_R U &= -\frac{\partial_R p}{\varrho} - \frac{m(R)}{R^2} \\ \partial_t \varrho &= -\frac{1}{R^2}\partial_R\left(R^2\varrho U\right). \end{aligned} \tag{18}$$

The stability analysis of the Bondi accretion in the Lagrange formulation has been performed by Balazs [13]. His approach is correct but inconclusive, because of a too stringent understanding of the notion of linearized stability. We show that using lagrangian variables one exactly reproduces the stability result already known from the Eulerian approach [7].

As the primary quantity one takes the deviation $\zeta(t,r) \equiv \Delta R(r,t)$ from the particle position in the unperturbed flow. The perturbed velocity reads $\Delta U = \partial_t^L\zeta \equiv (\partial_t + U\partial_R)\zeta$; here $\partial_t^L \equiv \partial_t + U\partial_R$ is the derivative with respect the Lagrangian time. The lagrangian perturbation $\Delta\varrho$ can be related to ζ by demanding that $\Delta m(R(r = \mathrm{const})) = 0$; that implies

$$\Delta\varrho = -\rho\left(\frac{2\zeta}{R} + \partial_R\zeta\right). \tag{19}$$

The U, a, ϱ solves Eqs. (18) with the mass function $m(R)$ including the contribution of the fluid itself. One derives from the first of Eqs. (18) the equation ruling ζ (take notice during the derivation that $\Delta m(R(r)) = 0$)

$$\begin{aligned} \left(\partial_t^L\right)^2\zeta &= \frac{2m(R)\zeta}{R^3} + \frac{1}{\varrho}\partial_R\left(a^2\varrho\left(\partial_R\zeta + \frac{2\zeta}{R}\right)\right) - \\ &\quad \frac{2\zeta}{\varrho}\partial_R p \end{aligned} \tag{20}$$

The perturbation of the other equation in (18) is satisfied identically.

Balazs tried to find solutions in the standard way, of the form $\zeta\left(R(r)\right), t) = \exp^{i\omega t}\zeta(R(r))$ with positive ω^2, that is with nonincreasing in time modulus; this clearly cannot be done, as shown below. The proper approach (adopted in the Eulerian variables by Moncrief [7]) is to define an energy of perturbed modes by the expression

$$\begin{aligned} E &= \int_V dV\varrho\left(\frac{1}{2}\left(\partial_t\zeta\right)^2 + \frac{1}{2}\left(\partial_R\zeta\right)^2\left(a^2 - U^2\right) + \right. \\ &\quad \left. \frac{\zeta^2}{R^2}\left(a^2 - \frac{m}{R} - R\partial_R a^2\right)\right), \end{aligned} \tag{21}$$

where V is an annulus between radii R_* and R_∞. The interesting and nontrivial question (since the last integrand in (21) can be negative in the vicinity of a sonic point) is whether the energy E of (21) is positive. It is enough to consider the region outside of the sonic horizon, because perturbations that exist inside will be washed away to the black hole.

In order to answer that question, notice an identity

$$\begin{aligned} \varrho\left(a^2 - \frac{m}{R} - R\partial_R a^2\right) &= -\partial_R\left[\varrho R\left(a^2 - \frac{m}{2R}\right)\right] + \\ \frac{2\varrho}{a^2 - U^2}\left(a^2 - \frac{m}{2R}\right)^2 &- 2\pi R^2\varrho; \end{aligned} \tag{22}$$

notice the occurence of the negative term at the end of the formula. Inserting (22) into the last term in the integrand of (21) and integrating by parts, one obtains after lengthy calculation that

$$E = \tilde{E} - \left[4\pi R\zeta^2\rho\left(a^2 - \frac{m}{2R}\right)\right]_{R_*}^{R_\infty} \tag{23}$$

where

$$\begin{aligned} \tilde{E} &= \frac{1}{2}\int_V dV\left(X^2 + Y^2\right) - \\ &\quad 2\pi\int_V dV\zeta^2\varrho^2, \end{aligned} \tag{24}$$

is not definite, while X and Y are given by

$$X = \sqrt{\varrho}\,\partial_t\zeta, \tag{25}$$

$$Y = \sqrt{\varrho}\left(\frac{2a^2R - m}{R^2\sqrt{a^2 - U^2}}\zeta + \sqrt{a^2 - U^2}\,\partial_R\zeta\right). \tag{26}$$

One can easily show that boundary terms of (23) are nonnegative. The occurence of a negative term in (24) means that the energy E of modes perturbing the selfgravitating flow may be negative.

The calculation of the derivative of the energy with respect to the Eulerian time yields

$$\begin{aligned} \partial_t E &= \int_V dV\zeta^2\frac{-\partial_t m(R)}{R^3} + \\ &\quad 4\pi\left[r^2\rho\left(\partial_t\zeta\partial_R\zeta\left(a^2 - U^2\right) - U\left(\partial_t\zeta\right)^2\right)\right]_{R_*}^{R_\infty}. \end{aligned} \tag{27}$$

A careful analysis of boundary terms shows that the contribution from R_∞ and R_* is negative definite. We can conclude that the outflow through the inner and outer boundaries is nonpositive and the energy E of perturbations cannot grow for the critical flow. Despite this fact, there is a possibility for unstable behaviour of the solutions of the time-dependent hydrodynamic equations, because the energy E is not positive definite.

The situation for subsonic flows is unclear even for test fluids, as pointed by Garlick [15]. While the analysis at R_∞ yields the same result, that the outflow diminishes the energy E, one has to assume that the inner outflow is completely absorbed by the surface of an attracting body, that is $\zeta(R_1, t) = 0$, in order to conclude that $\partial_t E \leq 0$. For more realistic cases, with the occurence of a reflection from the inner boundary, the energy of the perturbation may grow, signalling an instability, in addition to the ”bulk-type instability” suggested by the above analysis.

It is necessary to say in this place that the absence of exponentially growing linear modes does not mean that perturbed solutions of the nonlinear equation will be close to the background solution at all times. Linearized stability of a solution would imply only that evolving perturbations can be bounded by initial perturbations in a suitable sense. The existence of exponential linear modes in turn does not necessarily yields that perturbations grow indefinitely. Linear instability means that some evolving nonlinear perturbations are not bounded by initial perturbations; the ”strength” of evolving perturbations does not depend on the ”strength” of initial pertrbations [16].

Restricting our attention to test fluids, notice the following. The energy E is equal to $\tilde{E}$, and $\tilde{E}$ differs from the expression (24) only by the absence of the last, nonpositive integral, hence it is clearly positive. The time propagation in turn is like in (27), but without the integral $\zeta^2 \frac{-\partial_t m(R)}{R^3} = \zeta^2 \frac{4\pi U\rho}{R}$ on the right hand side. Thus E cannot grow, which excludes the exponential growth of X and Y and the long-term exponential growth of linear modes ζ. That does not guarantee that $|\zeta(R)|$ is time-independent.

This paper has been partially supported by the MNII grant 1PO3B 01229.

[1] H. Bondi, *Mon. Not. R. Astron. Soc.* **112**, 192(1952).
[2] F. C. Michel, *Astrophys. Space Sci.* **15**, 153(1972).
[3] S. Shapiro and S. Teukolsky, Black Holes, White Dwarfs and Neutron Stars, Wiley, New York, 1983.
[4] T.K. Das, *Astron. Astrophys.* **374**, 1150(2001); B. Kinasiewicz and T. Lanczewski, *Acta Phys. Pol.* **B36**, 1951(2005).
[5] J. Karkowski, B. Kinasiewicz, P. Mach E. Malec and Z. Świerczyński), *Phys. Rev.* **D73**, 021503(R)(2006).
[6] E. Malec, *Phys. Rev.* **D60**, 104043 (1999).
[7] V. Moncrief, *Astrophys. J.* **235**, 1038(1980).
[8] P. O. Mazur and E. Mottola (Los Alamos), *Proc. Nat. Acad. Sci.* **111**, 9545(2004).
[9] S. W. Hawking and G. F. R. Ellis, The Large Scale Structure of Space-Time, Cambridge University Press (1975).
[10] , A. H. Buchdahl, *Phys. Rev* **116**, 1027(1959).
[11] M. Abramowicz, W. Kluźniak and J.-P.Lassota , *Astronomy and Astrophysics*, **396**, L31(2002).
[12] A. E. Broderick and R. Narayan, *Astrophys.J.* **638**, L21(2006).
[13] N. L. Balazs, *Mon. Not. R. Astr. Soc.* **160**, 79(1972);
[14] E. Malec and N. O’Murchadha, *Phys. Rev.* **D50**, 6033(1994).
[15] A. R. Garlick, *Astr. Astrophysics*, **73**, 171(1979).
[16] G. Ioss and D. D. Joseph, Sec 2.7 in Elementary Stability and Bifurcation Theory, Springer Verlag New York Heildelberg Berlin 1980.